

Measure theoretic approach to the classification of cellular automata

Shin'ichirou Ishii

Kitakyushu National College of Technology, 5-20-1, Shii, Kokuraminamiku, Kitakyushu 803, Japan

Received 13 December 1989

Revised 20 June 1990

Abstract

Ishii, S., Measure theoretic approach to the classification of cellular automata, Discrete Applied Mathematics 39 (1992) 125–136.

Each measurable transformation that commutes with a shift on the set of configurations is proved to have an interesting dynamical property, i.e., for almost all initial configurations their orbits have a common structure. We can put those transformations into four classes according to their dynamical properties. The classification of cellular automata is derived from it.

1. Introduction

In [10], Wolfram observed that cellular automata can be put into four classes according to their dynamical behavior. A definition of these four classes based on the evolution on *finite configurations* was given by Culik II and Yu in [3]. In this paper we take another definition based on the fact that *for almost all initial configurations, their orbits have a common dynamical structure*. The author does not know what relations there are between the two ways of definition.

We introduce a class \mathcal{I} consisting of all measurable transformations which commute with at least one of the shifts on the set of configurations with a certain topology and a certain measure on it, and show that each element τ of \mathcal{I} has one of the following four dynamical properties.

- (1) For almost every initial configuration c_0 , its orbit $\tau^1(c_0), \tau^2(c_0), \dots$ evolves to a common homogeneous configuration in a bounded time.
- (2) For almost every initial configuration c_0 , its orbit $\tau^1(c_0), \tau^2(c_0), \dots$ evolves to an attractor which is cyclic in a sense in a bounded time.

Correspondence to: Professor S. Ishii, Department of Mathematics, Kyushu University 33, Fukuoka 812, Japan.

(3) For almost every initial configuration c_0 , its orbit $\tau^1(c_0), \tau^2(c_0), \dots$ evolves asymptotically to an attractor which is cyclic in a sense.

(4) For almost every initial configuration c_0 , its orbit $\tau^1(c_0), \tau^2(c_0), \dots$ evolves to no simple attractor.

The elements of \mathcal{S} can be put into four classes according to their dynamical properties described above. The classification of cellular automata is derived from that fact.

In Section 2 we define the topology and the measure on the set of configurations, in Section 3 the precise investigation of \mathcal{S} is done, in Section 4 we classify the set of cellular automata and in Section 5 we investigate the second class and the third class in detail.

2. Basic definitions and properties

In this paper, \mathbb{Z} denotes the set of integers, and n and q denote some fixed positive integers. We identify q with the set $\{0, 1, 2, \dots, q-1\}$ and call it a state set. The symbol \oplus denotes the addition of modulo q . Clearly $\langle q, \oplus \rangle$ is an Abelian group. For any i and j in \mathbb{Z} , $i \mid j$ denotes the fact that i is a divisor of j .

Definition 2.1. (1) A map $c: \mathbb{Z}^n \rightarrow q$ is called an (n, q) -*configuration*, or merely a *configuration*. \mathcal{C} denotes the set of all configurations.

(2) A configuration which is a constant map is called *homogeneous*. The symbol \bar{q}_0 denotes the constant map which takes the value q_0 .

We define a topological q -module and a measure on \mathcal{C} .

Definition 2.2. (1) \mathbb{T} denotes the product topology of the discrete topology on q , which is induced on $\mathcal{C} = q^{\mathbb{Z}^n}$.

(2) \mathbb{T}^2 denotes the product topology of \mathbb{T} , which is induced on $\mathcal{C} \times \mathcal{C}$.

Proposition 2.3. $\langle \mathcal{C}, \mathbb{T} \rangle$ and $\langle \mathcal{C} \times \mathcal{C}, \mathbb{T}^2 \rangle$ are metrizable, complete with such a metric.

Definition 2.4. The *addition* on \mathcal{C} is defined as follows:

$$(c_0 + c_1)(\alpha) = c_0(\alpha) \oplus c_1(\alpha),$$

for any $c_0, c_1 \in \mathcal{C}$ and $\alpha \in \mathbb{Z}^n$.

Proposition 2.5. $\langle \mathcal{C}, \mathbb{T}, + \rangle$ is a topological Abelian group.

Definition 2.6. (1) Let μ_0 be a probability measure on q defined as follows:

$$\mu_0(\{i\}) = \frac{1}{q}, \quad i = 0, \dots, q-1.$$

μ denotes its *product probability measure* defined on the field of Borel subsets of \mathcal{C} .

(2) Ω denotes the field of measurable subsets of \mathcal{C} with respect to the completion of μ .

(3) μ^2 denotes the product probability measure of μ on the field of Borel subsets of $\mathcal{C} \times \mathcal{C}$.

Definition 2.7. Let τ be a transformation on \mathcal{C} .

(1) We define τ to be *Borel measurable* if and only if for any Borel measurable subset \mathcal{B} of \mathcal{C} the set $\{c \in \mathcal{C} : \tau(c) \in \mathcal{B}\}$ is also Borel measurable.

(2) We define τ to be *measurable* if and only if for any measurable subset \mathcal{M} of \mathcal{C} the set $\{c \in \mathcal{C} : \tau(c) \in \mathcal{M}\}$ is also measurable.

Definition 2.8. Let τ_1 and τ_2 be two transformations on \mathcal{C} .

(1) We define the *addition* $\tau_1 + \tau_2$ and the *complement* $-\tau_1$ pointwisely.

(2) The *composition* $\tau_1 \circ \tau_2$ is defined as follows:

$$(\tau_1 \circ \tau_2)(c) = \tau_1(\tau_2(c)) \quad \text{for any } c \text{ in } \mathcal{C}.$$

The next proposition is easily derived from the definition.

Proposition 2.9. (1) Let τ_1 and τ_2 be two Borel measurable transformations on \mathcal{C} . The addition $\tau_1 + \tau_2$, the complement $-\tau_1$ and the composition $\tau_1 \circ \tau_2$ are Borel measurable.

(2) Each continuous transformation is Borel measurable.

Definition 2.10. Let $\beta \in \mathbb{Z}^n$. A *shift* s_β denotes the transformation on \mathcal{C} , which is defined as $s_\beta(c)(\alpha) = c(\alpha - \beta)$ for any $c \in \mathcal{C}$, $\alpha \in \mathbb{Z}^n$.

The next lemma is easily proved with the method of ergodic theory [8].

Lemma 2.11. For each $\beta \in \mathbb{Z}^n - \{0\}$:

(1) The shift s_β is continuous on $\langle \mathcal{C}, \mathbb{T} \rangle$.

(2) The shift s_β is ergodic on $\langle \Omega, \mathcal{C}, \mu \rangle$, i.e.,

(a) s_β is a measure preserving transformation on $\langle \Omega, \mathcal{C}, \mu \rangle$,

(b) if a subset A of \mathcal{C} is measurable and $s_\beta(A) = A$, then $\mu(A) = 0$ or 1.

The next lemma is derived from the continuity of shifts.

Lemma 2.12. Let c_1, c_2 be two elements of \mathcal{C} , $\tau_0, \tau_1, \tau_2, \dots$ a sequence of Borel measurable transformations on \mathcal{C} and β an element of \mathbb{Z}^n . If

$$\lim_{i \rightarrow \infty} \tau_i(c_1) = c_2$$

and

$$s_\beta \tau_i = \tau_i s_\beta$$

for any $i \geq 0$, then

$$\lim_{i \rightarrow \infty} \tau_i \circ s_\beta(c_1) = s_\beta(c_2).$$

Definition 2.13. Let c be an element of \mathcal{C} .

(1) A *subset* $\langle c \rangle$ of \mathcal{C} is defined as follows:

$$\langle c \rangle = \{s_\beta(c) \in \mathcal{C} : \beta \in \mathbb{Z}^n\}.$$

(2) Let c_1 and c_2 be two elements of \mathcal{C} . A binary relation \equiv on \mathcal{C} is defined as follows:

$$c_1 \equiv c_2 \text{ if and only if } \langle c_1 \rangle = \langle c_2 \rangle.$$

(3) The symbol \mathcal{C}/s denotes the set $\{\langle c \rangle : c \in \mathcal{C}\}$.

Definition 2.14. Let τ be a transformation on \mathcal{C} , c_0 an element of \mathcal{C} .

(1) For each nonnegative integer m , τ^m is defined inductively as follows:

For each $c \in \mathcal{C}$,

$$\tau^0(c) = c, \tag{1}$$

$$\tau^m(c) = \tau \circ \tau^{m-1}(c) \text{ for } m \geq 1. \tag{2}$$

(2) The set $\{\tau^m(c_0) : m \geq 1\}$ is called the *orbit* of τ from the initial configuration c_0 .

Definition 2.15. Let τ be a Borel measurable transformation on \mathcal{C} , T a nonnegative integer, t a positive integer, β an element of \mathbb{Z}^n and q_0 an element of \mathcal{Q} .

(1) $c \in \mathcal{C}$ is called (T, q_0) -stable if and only if the following property holds:

$$\tau^T(c) = \bar{q}_0, \tag{3}$$

$$\tau^{T'+1}(c) = \tau^{T'}(c) \text{ for any } T' \geq T. \tag{4}$$

$\mathcal{S}t_{T, q_0}^\tau$ denotes the set of all the (T, q_0) -stable configurations. We call \bar{q}_0 the limit point from c with the finite τ -iteration.

(2) $c \in \mathcal{C}$ is called (T, t, β) -cyclic if and only if the following property holds:

$$\tau^{T'+t}(c) = s_\beta \circ \tau^{T'}(c) \text{ for any } T' \geq T.$$

$\mathcal{C}y_{T, t, \beta}^\tau$ denotes the set of all the (T, t, β) -cyclic configurations. We call the set

$$\{\langle \tau^{T+i}(c) \rangle : 1 \leq i \leq t\}$$

the limit cycle from c with the finite τ -iteration. Note that the cardinal number of the limit cycle is a divisor of t .

(3) $c \in \mathcal{C}$ is called (T, t, β) -asymptotically cyclic if and only if the following property holds:

$$\lim_{T' \rightarrow \infty} (\tau^{T'}(c) - \tau^{T'}(c')) = \bar{0}$$

for some $c' \in \mathcal{C}y_{T,t,\beta}^\tau$.

$\mathcal{A}c_{T,t,\beta}^\tau$ denotes the set of all the (T, t, β) -asymptotically cyclic configurations. We call the set

$$\{\langle \tau^{T+i}(c') \rangle : 1 \leq i \leq t\}$$

the limit cycle from c with the infinite τ -iteration. Note that the cardinal number of the limit cycle is a divisor of t .

We can easily prove the following proposition from the definition above.

Proposition 2.16.

$$\mathcal{P}t_{T,q_0}^\tau \subset \mathcal{C}y_{T,1,0}^\tau, \quad (5)$$

$$\mathcal{C}y_{T,t,\beta}^\tau \subset \mathcal{A}c_{T,t,\beta}^\tau. \quad (6)$$

Lemma 2.17. $\mathcal{P}t_{T,q_0}^\tau$, $\mathcal{C}y_{T,t,\beta}^\tau$ and $\mathcal{A}c_{T,t,\beta}^\tau$ are measurable subsets of \mathcal{C} .

Proof. Clearly $\mathcal{P}t_{T,q_0}^\tau$ and $\mathcal{C}y_{T,t,\beta}^\tau$ are Borel subsets of \mathcal{C} . So they are measurable.

$\mathcal{A}c_{T,t,\beta}^\tau$ is proved to be a Σ_1^1 subset of \mathcal{C} from the definition. So the Souslin Theorem implies that it is measurable [2]. \square

Definition 2.18. (1) A subset \mathcal{J} of the set of Borel measurable transformations on \mathcal{C} is defined as follows. Let τ be a Borel measurable transformation.

$$\tau \in \mathcal{J} \quad \text{if and only if} \quad s_\alpha \tau = \tau s_\alpha \text{ for some } \alpha \in \mathbb{Z}^n - \{0\}.$$

(2) For τ in \mathcal{J} , a subset Γ_τ of \mathbb{Z}^n is defined as follows:

$$\Gamma_\tau = \{\alpha \in \mathbb{Z}^n - \{0\} : s_\alpha \tau = \tau s_\alpha\}.$$

3. Classification of \mathcal{J}

Definition 3.1. The four classes of Borel measurable transformations on \mathcal{C} , *CLASS I*, *CLASS II*, *CLASS III* and *CLASS IV* are defined as follows. Let τ be a Borel measurable transformation on \mathcal{C} .

(1) $\tau \in \text{CLASS I}$ if and only if the following property holds. For some $T \geq 0$ and $q_0 \in \mathcal{Q}$,

$$\mu(\mathcal{P}t_{T,q_0}^\tau) = 1.$$

(2) $\tau \in \text{CLASS II}$ if and only if the following property holds. For any $T \geq 0$ and $q_0 \in \mathcal{Q}$,

$$\mu(\mathcal{P}t_{T,q_0}^\tau) = 0,$$

and for some $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{C}y_{T,t,\beta}^\tau) = 1.$$

(3) $\tau \in \text{CLASS III}$ if and only if the following property holds. For any $T \geq 0$ and $q_0 \in q$,

$$\mu(\mathcal{P}t_{T,q_0}^\tau) = 0,$$

for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{C}y_{T,t,\beta}^\tau) = 0,$$

and for some $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{A}c_{T,t,\beta}^\tau) = 1.$$

(4) $\tau \in \text{CLASS IV}$ if and only if the following property holds. For any $T \geq 0$ and $q_0 \in q$,

$$\mu(\mathcal{P}t_{T,q_0}^\tau) = 0,$$

for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{C}y_{T,t,\beta}^\tau) = 0,$$

and for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{A}c_{T,t,\beta}^\tau) = 0.$$

Informally speaking, *CLASS I*, for example, is the class of all Borel measurable transformations such that the orbits of almost all initial configurations evolve to a common homogeneous fixed point in a bounded time. The next proposition is easily derived from the definition above.

Proposition 3.2. *Let $i, j = I, II, III$ or IV . If $i \neq j$,*

$$\text{CLASS } i \cap \text{CLASS } j = \emptyset.$$

Now we can state the classification theorem of \mathcal{I} .

Theorem 3.3. *Each element of \mathcal{I} belongs to one and only one of these four classes.*

The next lemma is essential to prove Theorem 3.3.

Lemma 3.4. *Let τ be an arbitrary element of \mathcal{I} .*

(1) *For any $T \geq 0$ and $q_0 \in q$,*

$$\mu(\mathcal{P}t_{T,q_0}^\tau) = 0 \text{ or } 1.$$

(2) *For any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,*

$$\mu(\mathcal{C}y_{T,t,\beta}^\tau) = 0 \text{ or } 1.$$

(3) For any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mu(\mathcal{A}c_{T,t,\beta}^{\tau}) = 0 \text{ or } 1.$$

Proof. We take one fixed element α of Γ_{τ} .

(1) Clearly the set $(\tau^T)^{-1}(\bar{q}_0)$ is invariant under s_{α} . So the ergodicity of the shift s_{α} implies the next equations:

$$\begin{aligned} \mu(\{c \in \mathcal{C}: \tau^T(c) = \bar{q}_0\}) &= \mu((\tau^T)^{-1}(\bar{q}_0)) \\ &= 0 \text{ or } 1. \end{aligned}$$

On the other hand, for any $T' \geq 0$ the set $(\tau^{T'+1} - s_{\beta} \circ \tau^{T'})^{-1}(\bar{0})$ is also invariant under s_{α} . Thus similarly we get the next relation. For any $T' \geq 0$

$$\begin{aligned} \mu(\{c \in \mathcal{C}: \tau^{T'+1}(c) = \tau^{T'}(c)\}) &= \mu((\tau^{T'+1} - \tau^{T'})^{-1}(\bar{0})) \\ &= 0 \text{ or } 1. \end{aligned}$$

Thus

$$\begin{aligned} \mu(\mathcal{P}t_{T,q_0}^{\tau}) &= \mu(\{c \in \mathcal{C}: \tau^T(c) = \bar{q}_0\} \cap \bigcup_{T' \geq T} \{c \in \mathcal{C}: \tau^{T'+1}(c) = \tau^{T'}(c)\}) \\ &= 0 \text{ or } 1. \end{aligned}$$

(2) Similarly we get the next relation. For any $T' \geq 0$

$$\begin{aligned} \mu(\{c \in \mathcal{C}: \tau^{T'+1}(c) = s_{\beta} \circ \tau^{T'}(c)\}) &= \mu((\tau^{T'+1} - s_{\beta} \circ \tau^{T'})^{-1}(\langle \bar{0} \rangle)) \\ &= 0 \text{ or } 1. \end{aligned}$$

Hence

$$\begin{aligned} \mu(\mathcal{C}y_{T,t,\beta}^{\tau}) &= \mu(\bigcup_{T' \geq T} \{c \in \mathcal{C}: \tau^{T'+1}(c) = s_{\beta} \circ \tau^{T'}(c)\}) \\ &= 0 \text{ or } 1. \end{aligned}$$

(3) Similarly from Lemma 2.12 we can prove that $\mathcal{A}c_{T,t,\beta}^{\tau}$ is invariant under s_{α} . Therefore

$$\mu(\mathcal{A}c_{T,t,\beta}^{\tau}) = 0 \text{ or } 1. \quad \square$$

Now we can prove Theorem 3.3.

Proof of Theorem 3.3. Let τ be an element of \mathcal{I} .

(1) If $\mu(\mathcal{P}t_{T,q_0}^{\tau}) = 1$ for some $T \geq 0$ and $q_0 \in q$, then $\tau \in \text{CLASS I}$.

(2) If $\mu(\mathcal{P}t_{T,q_0}^{\tau}) = 0$ for any $T \geq 0$ and $q_0 \in q$ and $\mu(\mathcal{C}y_{T,t,\beta}^{\tau}) = 1$ for some $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$, then $\tau \in \text{CLASS II}$.

(3) If $\mu(\mathcal{P}t_{T,q_0}^{\tau}) = 0$ for any $T \geq 0$ and $q_0 \in q$, $\mu(\mathcal{C}y_{T,t,\beta}^{\tau}) = 0$ for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$ and $\mu(\mathcal{A}c_{T,t,\beta}^{\tau}) = 1$ for some $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$, then $\tau \in \text{CLASS III}$.

(4) If $\mu(\mathcal{P}t_{T,q_0}^{\tau}) = 0$ for any $T \geq 0$ and $q_0 \in q$, $\mu(\mathcal{C}y_{T,t,\beta}^{\tau}) = 0$ for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$ and $\mu(\mathcal{A}c_{T,t,\beta}^{\tau}) = 0$ for any $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$, then $\tau \in \text{CLASS IV}$. \square

4. Classification of cellular automata

Definition 4.1. (1) Let r be a positive integer, $\eta_1, \eta_2, \dots, \eta_r$ pairwise distinct elements of \mathbb{Z}^n , $\mathcal{N} = \{\eta_1, \eta_2, \dots, \eta_r\}$ and σ a map from q^r into q . The pair $\langle \sigma, \mathcal{N} \rangle$ is called an *n-dimensional cellular automaton with q states*, or merely *CA*. The map σ and set \mathcal{N} are called respectively the *local map* and the *neighborhood indicator* of the CA.

(2) Let $\langle \sigma, \{\eta_1, \eta_2, \dots, \eta_r\} \rangle$ be a CA. The *global transformation* of a CA means the transformation τ on \mathcal{C} , which is defined as follows:

$$\tau(c)(\alpha) = \sigma(c(\alpha + \eta_1), \dots, c(\alpha + \eta_r)) \quad \text{for any } c \in \mathcal{C}, \alpha \in \mathbb{Z}^n.$$

We identify two CA's which have a common global transformation. So, each CA is denoted by its global transformation.

The next theorem is due to Hedlund [5].

Theorem 4.2. *Let τ be a continuous transformation on $\langle \mathcal{C}, \mathbb{T} \rangle$. τ is a CA if and only if $\tau \circ s_\alpha = s_\alpha \circ \tau$ for any $\alpha \in \mathbb{Z}^n$.*

Theorem 4.2 implies that each CA belongs to the class \mathcal{J} . Therefore the next theorem is derived from Theorem 3.3.

Theorem 4.3. *Each cellular automaton belongs to CLASS I, CLASS II, CLASS III or CLASS IV.*

The continuity of the cellular automaton τ implies that $\mathcal{P}_{T, q_0}^\tau$ and $\mathcal{C}_{T, t, \beta}^\tau$ are closed subsets of $\langle \mathcal{C}, \mathbb{T} \rangle$. The next proposition is easily derived from this fact.

Proposition 4.4. *Let τ be a CA.*

(1) *$\tau \in \text{CLASS I}$ if and only if the following property holds. For some $T \geq 0$ and $q_0 \in q$,*

$$\mathcal{P}_{T, q_0}^\tau = \mathcal{C}.$$

(2) *$\tau \in \text{CLASS II}$ if and only if the following property holds. For any $T \geq 0$ and $q_0 \in q$,*

$$\mathcal{P}_{T, q_0}^\tau \neq \mathcal{C},$$

and for some $T \geq 0$, $t \geq 1$ and $\beta \in \mathbb{Z}^n$,

$$\mathcal{C}_{T, t, \beta}^\tau = \mathcal{C}.$$

Remark. From the proposition above we get the following relations between Culik-Yu classes I, II and ours [3].

(1) If τ is in *CLASS I* in our sense, then τ is in *CLASS I* in Culik–Yu classification.

(2) If τ is in *CLASS I* \cup *CLASS II* in our sense, then τ is in *CLASS II* in Culik–Yu classification.

The author does not know whether the inverse relations hold or not. The relations on the higher classes are also unknown.

A presumptive correspondence between Wolfram’s classification and ours is in Table 1.

Table 1

The correspondence between Wolfram’s classification and ours

Wolfram’s classification	Our classification
CLASS I	<i>CLASS I</i>
CLASS II	<i>CLASS II</i>
CLASS III	<i>CLASS IV</i>
CLASS IV	<i>CLASS III</i>

According to the Wolfram’s observation [10], his *CLASS III* and *CLASS IV* are characterized as follows.

(1) τ is in *CLASS III* if and only if almost every evolution of τ leads to a chaotic pattern.

(2) τ is in *CLASS IV* if and only if almost every evolution of τ leads to complex localized structures, sometimes long lived.

Clearly, the character of our *CLASS IV* is nothing but that of his *CLASS III* described above. Indeed, we can prove that the CA of function code 42 [10] belongs to our *CLASS IV* from the fact that the shift invariant proper subgroup of $\langle \mathcal{C}, + \rangle$ has measure zero.

On the other hand, a close observation of the examples of Wolfram’s *CLASS IV*, function code 20 and 52 [10], show us that a stable or periodic configuration appears after all the “complex localized structures” died in a given space. The “complex localized structure” seems to be a phenomenon based on the way of convergence of point sequences in $\langle \mathcal{C}, \mathbb{T} \rangle$. That is the intuitive reason why we identified his *CLASS IV* as our *CLASS III*.

It is a difficult question to decide which classes in our sense a given CA belongs to. But from Proposition 4.4, we can say at least that *CLASS I* is recursive enumerable, that is, there is a Turing machine which halts if and only if we input a *CLASS I*-CA to it. Similarly, *CLASS I* \cup *CLASS II* is recursive enumerable.

5. On CLASS II and CLASS III

Definition 5.1. Let τ be a Borel measurable transformation on \mathcal{C} , T a nonnegative integer, t a positive integer, β an element of \mathbb{Z}^n and c_0, c_1, \dots, c_{t-1} be elements of \mathcal{C} such that $c_i \neq c_j$ when $i \neq j$.

(1) $c \in \mathcal{C}$ is called $(T, t, \beta; c_0, c_1, \dots, c_{t-1})$ -cyclic if and only if the following property holds:

$$\tau^{T+i}(c) \text{ is an element of } \langle c_i \rangle, \quad i=0, 1, \dots, t-1,$$

$$\tau^{T'+t}(c) = s_\beta \circ \tau^{T'}(c), \quad \text{for any } T' \geq T.$$

$\mathcal{C}y_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau$ denotes the set of all the $(T, t, \beta; c_0, c_1, \dots, c_{t-1})$ -cyclic configurations.

(2) $c \in \mathcal{C}$ is called $(T, t, \beta; c_0, c_1, \dots, c_{t-1})$ -asymptotically cyclic if and only if the following property holds: For some $c' \in \mathcal{C}y_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau$,

$$\lim_{T' \rightarrow \infty} (\tau^{T'}(c) - \tau^{T'}(c')) = \bar{0}.$$

$\mathcal{A}c_{T,t,\beta;c_0,\dots,c_{t-1}}^\tau$ denotes the set of all the $(T, t, \beta; c_0, \dots, c_{t-1})$ -asymptotically cyclic configurations.

We can easily prove the following proposition from the definition above.

Proposition 5.2.

$$\mathcal{C}y_{T,t,\beta}^\tau = \bigcup_{t' \mid t; t\beta' = t'\beta; \langle c_0 \rangle, \dots, \langle c_{t-1} \rangle \in \mathcal{C}/s} \mathcal{C}y_{T,t',\beta';c_0,c_1,\dots,c_{t'-1}}^\tau, \quad (7)$$

$$\mathcal{A}c_{T,t,\beta}^\tau = \bigcup_{t' \mid t; t\beta' = t'\beta; \langle c_0 \rangle, \dots, \langle c_{t-1} \rangle \in \mathcal{C}/s} \mathcal{A}y_{T,t',\beta';c_0,c_1,\dots,c_{t'-1}}^\tau. \quad (8)$$

We can prove the next lemma in the same way as the proof of Lemma 2.17.

Lemma 5.3. $\mathcal{C}y_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau$ and $\mathcal{A}c_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau$ are measurable subsets of \mathcal{C} .

Lemma 5.4. Let τ be arbitrary element of \mathcal{I} .

(1) For any $T \geq 0$, $t \geq 1$ and $c_0, c_1, \dots, c_{t-1} \in \mathcal{C}$

$$\mu(\mathcal{C}y_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau) = 0 \text{ or } 1.$$

(2) For any $T \geq 0$, $t \geq 1$ and $c_0, c_1, \dots, c_{t-1} \in \mathcal{C}$

$$\mu(\mathcal{A}c_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^\tau) = 0 \text{ or } 1.$$

Proof. We take one fixed element α of Γ_τ .

(1) For $i=0, 1, 2, \dots, t-1$, the set $(\tau^{T+i})^{-1}(\langle c_i \rangle)$ is invariant under s_α . So the ergodicity of s_α implies:

$$\begin{aligned}\mu(\{c \in \mathcal{C}: \tau^{T+i}(c) \in \langle c_i \rangle\}) &= \mu(\tau^{T+i})^{-1}(\langle c_i \rangle) \\ &= 0 \text{ or } 1.\end{aligned}$$

On the other hand, for any $T' \geq 0$

$$\begin{aligned}\mu(\{c \in \mathcal{C}: \tau^{T'+t}(c) = s_\beta \circ \tau^{T'}(c)\}) &= \mu((\tau^{T'+t} - s_\beta \circ \tau^{T'})^{-1}(\langle \bar{0} \rangle)) \\ &= 0 \text{ or } 1.\end{aligned}$$

Hence

$$\begin{aligned}\mu(\mathcal{E}y_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^T) &= \mu\left(\bigcup_{i=0,1,\dots,t-1} \{c \in \mathcal{C}: \tau^{T+i}(c) \in \langle c_i \rangle\} \right. \\ &\quad \left. \cap \bigcup_{T' \geq T} \{c \in \mathcal{C}: \tau^{T'+t}(c) = s_\beta \circ \tau^{T'}(c)\} \right) \\ &= 0 \text{ or } 1.\end{aligned}$$

(2) Similarly we can easily see that the set $\mathcal{A}c_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^T$ is invariant under s_α . Therefore

$$\mu(\mathcal{A}c_{T,t,\beta;c_0,c_1,\dots,c_{t-1}}^T) = 0 \text{ or } 1. \quad \square$$

Theorem 5.5.

- (1) For each τ in CLASS II one and only one of the following properties holds:
- (a) For almost all initial configurations their orbits evolve to a common limit cycle with the finite iteration.
 - (b) For almost every initial configuration c_1 and c_2 , their limit cycles with the finite iteration are distinct to each other.
- (2) For each τ in CLASS III one and only one of the following properties holds:
- (a) For almost all initial configurations their orbits evolve to a common limit cycle with the infinite iteration.
 - (b) For almost every initial configuration c_1 and c_2 , their limit cycles with the infinite iteration are distinct to each other.

Proof. (1) Let $\mu(\mathcal{E}y_{T,t,\beta}^T) = 1$. If for some $T' \leq T$, $t' \mid t$, $\beta' = (t'/t)\beta$ and $\langle c_0 \rangle, \dots, \langle c_{t'-1} \rangle \in \mathcal{C}/s$,

$$\mu(\mathcal{E}y_{T',t',\beta';c_0,c_1,\dots,c_{t'-1}}^{T'}) = 1.$$

Then clearly (a) holds. So we can assume the following fact. For any $T' \leq T$, $t' \mid t$, $\beta' = (t'/t)\beta$ and $\langle c_0 \rangle, \dots, \langle c_{t'-1} \rangle \in \mathcal{C}/s$,

$$\mu(\mathcal{E}y_{T',t',\beta';c_0,c_1,\dots,c_{t'-1}}^{T'}) = 0.$$

Then for any $T' \leq T$, $t' \mid t$, $\beta' = (t'/t)\beta$ and $\langle c_0 \rangle, \dots, \langle c_{t'-1} \rangle \in \mathcal{C}/s$,

$$\mu^2((\mathcal{E}y_{T',t',\beta';c_0,c_1,\dots,c_{t'-1}}^{T'})^2) = 0.$$

On the other hand

$$\mu^2((\mathcal{C}y_{T,t,\beta}^{\tau})^2) = 1.$$

Therefore (b) holds.

(2) It is proved similarly as above. \square

Acknowledgement

The author wishes to express his heartfelt thanks to Professor S. Huzino who advised and encouraged him to make this work.

References

- [1] H. Aso and N. Honda, Dynamical characteristics of linear cellular automata, *J. Comput. System Sci.* 30 (1985) 219–317.
- [2] J. Barwise, *Handbook of Mathematical Logic* (North-Holland, Amsterdam, 1985).
- [3] K. Culik II and S. Yu, Undecidability of CA classification schemes, *Complex Systems* 2 (1988) 177–190.
- [4] F. Fogelman, Y. Robert and M. Tchuente, *Automata Networks in Computer Science* (Manchester University Press, Manchester, CT, 1987).
- [5] G.A. Hedlund, Endomorphisms and automorphisms of the shift dynamical system, *Math. Systems Theory* 3 (1969) 320–375.
- [6] P.G. Hinman, *Recursion-Theoretic Hierarchies* (Springer, Berlin, 1978).
- [7] F. Robert, *Discrete Iterations* (Springer, Berlin, 1986).
- [8] P. Walters, *An Introduction to Ergodic Theory* (Springer, Berlin, 1982).
- [9] J.S. Willson, On the ergodic theory of cellular automata, *Math. Systems Theory* 9 (1975) 132–141.
- [10] S. Wolfram, Universality and computability in cellular automata, *Phys. D* 10 (1984) 1–35.